



24730 Avenue 13
Madera, CA 93637
Phone: (559) 661-6386
FAX: (559) 661-6135

PLANT TISSUE Sampling Guidelines

DECIDUOUS FRUIT

APPLE	OLIVE
APRICOT	PEACH
ASIAN PEAR	PEAR
CHERRY	PLUM
FIG	PRUNE
NECTARINE	

Recommended timing

Leaf tissue samples may be collected at any time during the growing season. For many elements, the least change in concentration occurs during late June to July. University of California guidelines are usually correlated to leaf tissue samples taken in July. Interpretation of tissue samples taken at times other than July are generally interpolated from a knowledge of how nutrient concentration changes as the season progresses and from historical database information.

Sampling

For all leaf samples, select trees which are of similar age, variety, rootstock, and vigor. Leaves should be taken about 4-6 feet above ground. Always sample from the current season non-fruiting shoots or spurs. Select a leaf from each of the four quadrants of an individual tree, and sample 20-25 trees from the block. Thus, the sample should be a composite of about 80-100 leaves.

Apples
Apricots
Asian pears
Cherries
Pear
Plums
Prunes

March to April..... Select the most recently matured leaf from the base of the spur (usually the basal first to third leaves).
May Select the most recently matured leaf from the tip of the spur (usually the terminal first to third leaves).
June to October Select the terminal leaf on the spur.
Hull samples for boron..... Collect hulls at 4-6 feet above ground from a number of trees at harvest. The total sample should have about 40-50 hulls.

Figs
Nectarines
Olives
Peaches

April to October Select basal to mid-shoot leaves.


Preferred analyses

Whole leaves – Test code 1254.

PLANT TISSUE Sampling Guidelines

DECIDUOUS FRUIT

APPLES
APRICOTS
ASIAN PEARS
CHERRIES
PEARS
PLUMS
PRUNES





24730 Avenue 13
Madera, CA 93637
Phone: (559) 661-6386
FAX: (559) 661-6135

PLANT TISSUE Sampling Guidelines

DECIDUOUS FRUIT

FIGS
NECTARINES
OLIVES
PEACHES

